

10/536613

JC13 Rec'd PCT/PTO 25 MAY 2005

**SINOMENINE AND SINOMENINE COMPOUNDS**

---

**SHANGHAI INSTITUTE OF MATERIA MEDICA,  
Chinese Academy of Sciences  
Zu chong-zhi Road 555  
Zhangjiang Hi-Tech Park  
Pudong - SHANGHAI  
PEOPLE'S REPUBLIC OF CHINA**

**LES LABORATOIRES SERVIER  
12, Place de La Défense  
92415 COURBEVOIE Cedex  
FRANCE**

**INVENTORS :**

<b>QIN Guo-Wei</b>
<b>TANG Xi-Can</b>
<b>WANG Rui</b>
<b>ZHOU Tian-Xi</b>
<b>LESTAGE Pierre</b>
<b>CAIGNARD Daniel-Henri</b>
<b>RENARD Pierre</b>

**Title of the invention :**

Sinomenine and sinomenine compounds

**Field of the invention :**

The invention relates to sinomenine and sinomenine derivatives and their use in  
5 mnemocognitive disorders.

**Background and Prior art of the invention :**

*Sinomenium acutum* is a plant taking the form of a ligneous liana, which is widespread in the centre, South-East and South-West of China and is included in the Chinese Pharmacopoeia (Pharmacopoeia Committee of People's Republic of China, 2000). It  
10 contains a large number of alkaloids of various chemical structures, such as sinomenine, sinoacutine, ethylsinomenine, disinomenine, tetrahydroepiberberine, tuduranine and magnoflorine (Huang Tai-Kang, « *Handbook of the Composition and Pharmacology of Common Chinese Drugs* », Chinese Medical Science and Technology Publisher, 1994, Beijing, 1156-1160).

15 Sinomenine, a morphine-like alkaloid and a major constituent of the plant, has been much studied; in particular, it has been possible to demonstrate anti-inflammatory, immunosuppressive, anti-arrhythmic and analgesic properties (Qiang Liu *et al.*, Chinese Traditional and Herbal Drugs, 1997, 28(4), 247).

We have now discovered that sinomenine has mnemocognition-facilitating properties in  
20 animal experimental models.


Ageing of the population due to increased life expectancy has brought with it a major increase in cognitive disorders associated with normal cerebral ageing or pathological cerebral ageing occurring in the course of neurodegenerative diseases such as, for example, Alzheimer's disease.

25 The majority of substances used today in treating cognitive disorders associated with ageing act by facilitating the central cholinergic systems – either directly, as in the case of acetylcholinesterase inhibitors (tacrine, donepezil) and cholinergic agonists (nefiracetam),

or indirectly, as in the case of nootropic agents (piracetam, pramiracetam) and cerebral vasodilators (vinpocetine).

It has been therefore been especially valuable to synthesise new compounds that are capable of opposing the cognitive disorders associated with ageing and/or of improving cognitive processes.


The present invention relates, on the one hand, to the use of sinomenine :



and/or sinomenine compounds in mnemocognitive disorders and, on the other hand, to the synthesis of new compounds having especially valuable pharmacological properties in the same area.

#### **Detailed description of the invention :**

The present invention relates more specifically to compounds of formula (I) :



wherein

- R<sub>1</sub> represents an alkyl group,
- R<sub>2</sub> represents a hydrogen atom or an alkylcarbonyl group, an haloalkylcarbonyl group or an arylcarbonyl group,

- Y represents a group  $\text{NR}_7$ ,  $\text{N}^+\text{O}^-\text{R}_7$  or  $\text{N}^+\text{R}_7\text{R}'_7 \text{Z}^-$

wherein  $\text{R}_7$  and  $\text{R}'_7$ , identical or different, each represent an alkyl group, and  $\text{Z}^-$  represents a halogen anion,

- $\text{R}_3$  represents a hydroxy or alkoxy group,
- $\text{R}_4$  and  $\text{R}'_4$  each represent a hydrogen atom or together form an additional bond, or  $\text{R}_3$  and  $\text{R}_4$  together form an oxo or  $=\text{N}-\text{OR}_8$  group (wherein  $\text{R}_8$  represents a hydrogen atom or an alkyl group),
- $\text{R}_6$  represents a hydroxy, alkylcarbonyloxy (wherein the alkyl moiety can be substituted by a hydroxy, alkoxy, carboxy or alkyloxycarbonyl group) or alkoxy group,
- $\text{R}_5$  and  $\text{R}'_5$  each represent a hydrogen atom or together form an additional bond, or  $\text{R}_5$  and  $\text{R}_6$  together form an oxo,  $=\text{N}-\text{OR}_9$  or  $=\text{N}-\text{NR}_{10}\text{R}_{11}$  group (wherein  $\text{R}_9$ ,  $\text{R}_{10}$ , and  $\text{R}_{11}$ , which may be the same or different, each represent a hydrogen atom or an alkyl group),
- and X represents a halogen atom,

with the proviso that the compound of formula (I) cannot represent 1-bromo-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one,

it being understood that

- "alkyl" means an alkyl group containing 1 to 6 carbon atoms which may be linear or branched,
- "alkoxy" means an alkyloxy group containing 1 to 6 carbon atoms which may be linear or branched,


to their enantiomers and diastereoisomers, and to addition salts thereof with a pharmaceutically acceptable acid or base.

Among the pharmaceutically acceptable acids there may be mentioned, without implying any limitation, hydrochloric acid, hydrobromic acid, sulphuric acid, phosphonic acid,

acetic acid, trifluoroacetic acid, lactic acid, pyruvic acid, malonic acid, succinic acid, glutaric acid, fumaric acid, tartaric acid, maleic acid, citric acid, ascorbic acid, methanesulphonic acid, camphoric acid, oxalic acid etc..

Among the pharmaceutically acceptable bases there may be mentioned, without implying  
5 any limitation, sodium hydroxide, potassium hydroxide, triethylamine, tert-butylamine etc..

The preferred configuration of compounds of formula (I) according to the invention is that shown in formula (I'):



The preferred group  $R_1$  is the methyl group.

10 Advantageously,  $R_2$  represents a hydrogen atom or a group  $\text{EtCO}$  and more preferably a hydrogen atom.

Y represents, preferably, a group  $\text{NR}_7$  or  $\text{N}^+\text{O}^-\text{R}_7$  and, more especially, a group  $\text{N}^+\text{Me}$   
or  $\text{N}^+\text{MeO}^-$ .

X represents, very preferably, a chlorine or bromine atom.


15 Advantageously, the invention relates to compounds of formula (I) wherein  $R_3$  represents an alkoxy group and  $R_4$  and  $R'_4$  together form an additional bond.

The preferred meaning of  $R_5$  is a hydrogen atom.

$R_6$  represents advantageously an OH, ethoxy or alkylcarbonyloxy group and, more especially, ethylcarbonyloxy.

Another interesting aspect of the invention is compounds of formula (I) for which  $R_5$  and  $R_6$  form together an oxo group or an  $=N-OH$  group.

Very preferably, the invention relates to compounds of formula (I'') and (I''') :




5      wherein  $Y'$  represents  $\text{>N-Me}$  or  $\text{>N}^+\text{O}^-$ ,  $R'_2$  and  $R'_6$ , which may be the same or different, represent a hydrogen atom or an alkylcarbonyl group,  $X'$  represents a chlorine or bromine atom and  $Z$  represents  $=O$  or  $=N-OH$ .


Even more preferably, the invention relates to compounds of formula (I) that are (9 $\alpha$ ,13 $\alpha$ )-1-chloro-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-4,6-diol, (9 $\alpha$ ,13 $\alpha$ )-1-chloro-3,7-dimethoxy-17-methyl-4-(propionyloxy)-7,8-didehydromorphinan-6-yl propionate, (9 $\alpha$ ,13 $\alpha$ )-1-bromo-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-4,6-diol, (9 $\alpha$ ,13 $\alpha$ )-1-bromo-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one oxime, (9 $\alpha$ ,13 $\alpha$ )-1-bromo-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one N-oxide, (9 $\alpha$ ,13 $\alpha$ )-1-chloro-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one N-oxide.

The enantiomers and diastereoisomers and addition salts with a pharmaceutically acceptable acid or base of the preferred compounds of the invention form an integral part of the invention.

The invention relates also to a process for the preparation of compounds of formula (I), which process is characterised in that there is used as starting material the compound of formula (II) :




- 5      obtained by extraction starting from the stem of *Sinomenum acutum* according to Figure 1 below :



**Figure 1** : Extraction of the compound of formula (II)



which is subjected to the action of a halogenating agent such as  $\text{SO}_2\text{Cl}_2$  or  $\text{Br}_2$  to obtain the compound of formula (I/a), a particular case of the compounds of formula (I):



wherein X is as defined for formula (I), which compound of formula (I/a) is subjected to conventional chemical reactions to obtain the totality of the compounds of formula (I), which may be purified according to a conventional separation technique, are converted, if desired, into their addition salts with a pharmaceutically acceptable acid or base and are separated, where appropriate, into their isomers according to a conventional separation technique.

Besides the fact that the compounds of the present invention are new, they possess properties of facilitating cognitive processes, making them of use in the treatment of cognitive deficiencies associated with cerebral ageing and with neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, Pick's disease, Korsakoff's disease, and frontal lobe and subcortical dementias.


The invention relates also to pharmaceutical compositions comprising as active ingredient at least one compound of formula (I) together with one or more appropriate, inert, non-toxic excipients.

The Applicant has moreover discovered that sinomenine and/or sinomenine compounds have mnemocognition-facilitating properties.

The invention accordingly relates also to the use of sinomenine and/or sinomenine compounds in obtaining pharmaceutical compositions for use in the treatment of cognitive

deficiencies associated with cerebral ageing and with neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, Pick's disease, Korsakoff's disease, and frontal lobe and subcortical dementias.

More especially, the invention relates to the use, in obtaining pharmaceutical compositions for use in the treatment of cognitive deficiencies associated with cerebral ageing and with neurodegenerative diseases, of sinomenine and/or sinomenine compounds such as, for example, the compounds of formula (Ia) :



wherein  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R'_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$  and  $Y$  are as defined for formula (I), and, more especially, of (9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one hydrazone; (7 $\alpha$ ,9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methylmorphinan-6-one; (7 $\beta$ ,9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methylmorphinan-6-one; (9 $\alpha$ ,13 $\alpha$ )-3,7-dimethoxy-17-methyl-6-oxo-7,8-didehydromorphinan-4-yl propionate; (9 $\alpha$ ,13 $\alpha$ )-3,4,7-trimethoxy-17-methyl-7,8-didehydromorphinan-6-one; (9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one oxime; (9 $\alpha$ ,13 $\alpha$ )-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-4,6-diol; (9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one N-oxide; (9 $\alpha$ ,13 $\alpha$ )-6-amino-3,7-dimethoxy-17-methylmorphinan-4-ol; 4-[(9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-yl]-oxy}-4-oxobutanoic acid; (9 $\alpha$ ,13 $\alpha$ )-3,7-dimethoxy-17-methyl-4-(propionyloxy)-7,8-didehydromorphinan-6-yl propionate; (9 $\alpha$ ,13 $\alpha$ )-17-benzyl-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-17-ium-6-one bromide; (9 $\alpha$ ,13 $\alpha$ )-17-ethyl-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-17-ium-4,6-diol bromide; (9 $\alpha$ ,13 $\alpha$ )-17-ethyl-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-17-ium-6-one

bromide; (9 $\alpha$ ,13 $\alpha$ )-4-(benzoyloxy)-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-yl benzoate; (9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-yl benzoate; (9 $\alpha$ ,13 $\alpha$ )-6-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-4-yl benzoate.

- 5 An advantageous aspect of the invention relates to the use of sinomenine in obtaining pharmaceutical compositions for use in the treatment of cognitive deficiencies associated with cerebral ageing and with neurodegenerative diseases.

Another especially interesting aspect of the invention relates to the use, in obtaining pharmaceutical compositions for use in the treatment of cognitive deficiencies associated with cerebral ageing and with neurodegenerative diseases, of compounds of formula (Ia)

10 and, more especially, of (9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one hydrazone; of (7 $\alpha$ ,9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methylmorphinan-6-one; of (7 $\beta$ ,9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methylmorphinan-6-one; of (9 $\alpha$ ,13 $\alpha$ )-3,7-dimethoxy-17-methyl-6-oxo-7,8-didehydromorphinan-4-yl propionate; of

15 (9 $\alpha$ ,13 $\alpha$ )-3,4,7-trimethoxy-17-methyl-7,8-didehydromorphinan-6-one; of (9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one oxime; of (9 $\alpha$ ,13 $\alpha$ )-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-4,6-diol; of (9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one N-oxide; of (9 $\alpha$ ,13 $\alpha$ )-6-amino-3,7-dimethoxy-17-methylmorphinan-4-ol; of 4-[(9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-

20 methyl-7,8-didehydromorphinan-6-yl]oxy}-4-oxobutanoic acid; of (9 $\alpha$ ,13 $\alpha$ )-3,7-dimethoxy-17-methyl-4-(propionyloxy)-7,8-didehydromorphinan-6-yl propionate; of (9 $\alpha$ ,13 $\alpha$ )-17-benzyl-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-17-ium-6-one bromide; of (9 $\alpha$ ,13 $\alpha$ )-17-ethyl-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-17-

ium-4,6-diol bromide; of (9 $\alpha$ ,13 $\alpha$ )-17-ethyl-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-17-ium-6-one bromide; of (9 $\alpha$ ,13 $\alpha$ )-4-(benzoyloxy)-3,7-dimethoxy-

25 17-methyl-7,8-didehydromorphinan-6-yl benzoate; of (9 $\alpha$ ,13 $\alpha$ )-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-yl benzoate; of (9 $\alpha$ ,13 $\alpha$ )-6-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-4-yl benzoate.


The invention relates also to pharmaceutical compositions comprising sinomenine or a compound thereof, in combination with one or more pharmaceutically acceptable excipients, for use in the treatment of cognitive deficiencies associated with cerebral ageing and with neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, Pick's disease, Korsakoff's disease, and frontal lobe and subcortical dementias.

Among the pharmaceutical compositions according to the invention, there may be mentioned more especially those that are suitable for oral, parenteral (intravenous or subcutaneous) or nasal administration, tablets or dragées, sublingual tablets, gelatin capsules, lozenges, suppositories, creams, ointments, dermal gels, injectable preparations, drinkable suspensions etc..

The useful dosage can be varied according to the nature and severity of the disorder, the administration route and also the age and weight of the patient. The dosage varies from 0.01 mg to 1 g per day in one or more administrations.

The following Examples illustrate the invention but do not limit it in any way.

**EXAMPLE 1 : (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one**



To a solution of 100 mg of the compound of formula (II) in 5 ml of  $\text{CHCl}_3$  there are added 3 drops of  $\text{SO}_2\text{Cl}_2$ . The reaction mixture is stirred at ambient temperature for 4 hours and

the pH is adjusted to 7-8 with NaHCO<sub>3</sub> solution; extraction with CHCl<sub>3</sub> is then carried out. The organic phase is evaporated under reduced pressure and the residue obtained is chromatographed on silica gel using an eluant CHCl<sub>3</sub>-MeOH (9 : 1) to yield the title compound in the form of a yellowish solid.

5 *Melting point* : 126-128°C.

**EXAMPLE 2 : (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-3,7-dimethoxy-17-methyl-6-oxo-7,8-didehydro-morphinan-4-yl propionate**

To a solution of 500 mg of the compound obtained in Example 1 and 100 mg of DMAP in 15 ml of pyridine there are slowly added 2 ml of propionic anhydride, and the reaction  
10 mixture is stirred at ambient temperature for 3 hours. The reaction mixture is then evaporated and the residue obtained is dissolved in a small volume of water. The solution obtained is adjusted to pH = 8-9 with NaHCO<sub>3</sub> solution and is then extracted with CHCl<sub>3</sub>. The organic phase is washed 3 times with water, dried over sodium sulphate and evaporated. The residue obtained is chromatographed on silica gel using an eluant CHCl<sub>3</sub>-MeOH  
15 (20 : 1) to yield the title compound in the form of a colourless oil.

**EXAMPLE 3 : (6 $\beta$ ,7 $\beta$ ,9 $\alpha$ ,13 $\alpha$ )-1-Chloro-3,7-dimethoxy-17-methylmorphinan-4,6-diol**

A mixture of 720 mg of compound of Example 1 and 100 mg of PtO<sub>2</sub> in 50 ml of absolute ethanol is stirred at room temperature under H<sub>2</sub> atmosphere for 12 hours. The PtO<sub>2</sub> is  
20 removed by filtration and the ethanol is evaporated in vacuum to give a syrupy residue. This residue is washed with hot absolute ethanol (10 ml) to give a powdery solid which is collected by filtration and crystallized in CHCl<sub>3</sub>/C<sub>2</sub>H<sub>5</sub>OH to give the title compound in the form of white crystals.

*Melting point* : 210-212°C.

**EXAMPLE 4 : (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-3,7-dimethoxy-17-methyl-7,8-didehydro-morphinan-4,6-diol**

To a solution of 500 mg of the compound obtained in Example 1 in 15 ml of methanol there are added 500 mg of NaBH<sub>4</sub>, and the reaction mixture is stirred for 1.5 hours. The methanol is then evaporated off and the residue obtained is extracted with CHCl<sub>3</sub>. The organic phase is dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated under reduced pressure. The title compound is obtained in the form of white crystals, by recrystallisation from Et<sub>2</sub>O.

*Melting point* : 118-120°C.

**EXAMPLE 5 : (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-3,7-dimethoxy-17-methyl-4-(propionyloxy)-7,8-didehydromorphinan-6-yl propionate**

The title compound is obtained using the procedure described for Example 2, starting from the compound obtained in Example 4.

Oil.

**EXAMPLE 6 : (6 $\beta$ ,7 $\beta$ ,9 $\alpha$ ,13 $\alpha$ )-1-Chloro-3,7-dimethoxy-17-methyl-4-(propionyloxy)-morphinan-6-yl propionate**

The title compound is obtained using the procedure described for Example 2, starting from the compound obtained in Example 3.

Oil.

**EXAMPLE 7 : (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-3,4,7-triméthoxy-17-methyl-7,8-didehydro-morphinan-6-one**

A solution of 400 mg of the compound of Example 1 in 10 ml of methanol is treated with an excess of a freshly made preparation of diazomethane in ether, and the reaction mixture is stirred at ambient temperature for 12 hours. The excess of diazomethane is then broken down using glacial acetic acid, and the solvents are evaporated off under reduced pressure.

The residue obtained is adjusted to pH = 8-9 using saturated NaHCO<sub>3</sub> solution and is then

extracted with  $\text{CHCl}_3$ . The organic phase is dried over  $\text{Na}_2\text{SO}_4$  and evaporated *in vacuo*, and the residue obtained is chromatographed on silica gel (eluant  $\text{CHCl}_3$ -MeOH) to yield the title product in the form of an oil.

**EXAMPLE 8 : (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-3,4,7-trimethoxy-17-méthyl-7,8-didehydro-morphinan-6-ol**

The title compound is obtained using the procedure described for Example 4, starting from the compound obtained in Example 7. Colourless crystals.

*Melting point* : 163-165°C.

**EXAMPLE 9 : (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one oxime**

To a solution of 360 mg of the compound obtained in Example 1 in ethanol there are added 200 mg of  $\text{NH}_2\text{OH} \cdot \text{HCl}$  and 300 mg of sodium acetate. The reaction mixture is stirred for 4 hours; it is then filtered and evaporated under reduced pressure. The residue obtained is made alkaline using  $\text{NaHCO}_3$  solution and extracted with  $\text{CHCl}_3$ . The organic phase is dried over  $\text{Na}_2\text{SO}_4$  and then evaporated under reduced pressure, and the title compound is obtained in the form of needles, by recrystallisation from EtOH.

*Melting point* : 167-169°C.

**EXAMPLE 10 : (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-6-ethoxy-4-hydroxy-3-methoxy-17-methyl-5,6-didehydromorphinan-7-one**

To a solution of 1.3 g of the compound obtained in Example 1 in 100 ml of  $\text{CHCl}_3$  and 10 ml of absolute alcohol there is added  $\text{SO}_2\text{Cl}_2$  at 10°C, and the reaction mixture is stirred for 8 hours. The solvent is then evaporated off under reduced pressure, and the residue is neutralised using  $\text{NaHCO}_3$  and then extracted with  $\text{CHCl}_3$ . The organic phase is dried over  $\text{Na}_2\text{SO}_4$  and evaporated under reduced pressure, and the title compound is obtained in the form of yellowish crystals, by recrystallisation from  $\text{CH}_3\text{CN}$ .

*Melting point* : 190-192°C.

**EXAMPLE 11 : (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-4-hydroxy-3,7-dimethoxy-17-méthyl-7,8-didehydromorphinan-6-one hydrazone**

A solution of 600 mg of the compound obtained in Example 1 in 10 ml of 85 % hydrazine is stirred at 90°C for 8 hours. After cooling, the reaction mixture is filtered and the solid  
5 obtained is washed with water and recrystallised from EtOH to yield the title compound in the form of yellowish crystals.

*Melting point* : 235-237°C.

**EXAMPLE 12 : (9 $\alpha$ ,13 $\alpha$ )-1-Bromo-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one**

A solution of 6.6 g of the compound obtained in Example 1 in 150 ml of CHCl<sub>3</sub> is cooled to 0°C, and bromine dried over concentrated sulphuric acid is added dropwise, with stirring, whilst maintaining the reaction mixture at 5°C. The reaction is continued for a few minutes and then neutralisation is carried out using NaHCO<sub>3</sub>. The organic phase is separated off, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated under reduced pressure, and the residue  
10 obtained is recrystallised from EtOH to yield the title compound in the form of brown crystals.

*Melting point* : 163-165°C.

**EXAMPLE 13 : (9 $\alpha$ ,13 $\alpha$ )-1-Bromo-3,7-dimethoxy-17-methyl-7,8-didehydro-morphinan-4,6-diol**

The title compound is obtained using the procedure described for Example 4, starting from the compound obtained in Example 12 and replacing NaBH<sub>4</sub> by KBH<sub>4</sub>.

Solid.

*Melting point* : 144-146°C.



**EXAMPLE 14 : (9 $\alpha$ ,13 $\alpha$ )-1-Bromo-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one hydrazone**

The title compound is obtained using the procedure described for Example 11, starting from the compound obtained in Example 12.

5 Solid.

*Melting point* : 208-210°C.

**EXAMPLE 15 : (9 $\alpha$ ,13 $\alpha$ )-1-Bromo-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one oxime**

10 The title compound is obtained using the procedure described for Example 9, starting from the compound obtained in Example 12.

Solid.

*Melting point* : 180-182°C.

**EXAMPLE 16 : (9 $\alpha$ ,13 $\alpha$ )-1-Bromo-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one N-oxide**

15 A mixture of compound of Example 12 (820 mg) in H<sub>2</sub>O<sub>2</sub> (10 ml) is stirred at room temperature for 24 hours and then extracted with CHCl<sub>3</sub> three times (30 ml X 3). The combined extracts are dried overnight with anhydrous Na<sub>2</sub>SO<sub>4</sub> and the solvent is removed by evaporation to give a residue to which are added 30 ml of cold water. The powdery solid is collected by filtration, washed with cold water until the water being colorless, and  
20 crystallized in ethanol to give the title compound as a solid.

*Melting point* : 170-172°C.

**EXAMPLE 17 : (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-6-one N-oxide**

25 A mixture of compound of Example 1 (720 mg) in H<sub>2</sub>O<sub>2</sub> (10 ml) is stirred at room temperature for 24 hours and then extracted three times (25 ml X 3). The combined

extracts are dried over anhydrous  $\text{Na}_2\text{SO}_4$  and the solvent is removed by evaporation. 30 ml of cold water are added to the residue obtained and the resulting powdery white solid is collected by filtration and crystallized in ethanol to give the title compound as a solid.

*Melting point* : 170-172°C.

5     **EXAMPLE 18 : (9 $\alpha$ ,13 $\alpha$ )-1-Bromo-3,7-dimethoxy-17-methylmorphinan-4,6-diol**

The title compound is obtained using the procedure described for Example 3, starting from the compound obtained in Example 12.

*Melting point* : 160-162°C.

10     **EXAMPLE 19 : (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-6-ethoxy-3-methoxy-17-methyl-5,6-didehydromorphinan-4,7-diol**

The title compound is obtained using the procedure described for Example 4, starting from the compound obtained in Example 10 and replacing  $\text{NaBH}_4$  by  $\text{KBH}_4$ .

Solid.

*Melting point* : 168-170°C.

15     **EXAMPLE 20 : (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-6-ethoxy-4-hydroxy-3-methoxy-17-methyl-5,6-didehydromorphinan-7-one oxime**

The title compound is obtained using the procedure described for Example 9, starting from the compound obtained in Example 10.

Solid.


20     *Melting point* : 216-218°C.

**EXAMPLE 21 : (9 $\alpha$ ,13 $\alpha$ )-17-Benzyl-1-bromo-4-hydroxy-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-17-ium-6-one bromide**

The title compound is obtained from compound of example 12 subjected to the action of benzylbromide.

Melting point : 190-192°C.

**EXAMPLE 22 :**



The title compound is obtained after treatment of compound of Example 1 in basic medium.

5 Melting point : 204-206°C.

**EXAMPLE 23 :** (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-4-hydroxy-6-isopropoxy-3-methoxy-17-methyl-5,6-didehydromorphinan-7-one

The title compound is obtained using the procedure described in Example 10 replacing absolute alcohol with isopropyl alcohol.

10 Melting point : 216-218°C.

**EXAMPLE 24 :** (9 $\alpha$ ,13 $\alpha$ )-1-Chloro-3,7-dimethoxy-17-methyl-6-oxo-7,8-didehydromorphinan-4-yl chloroacetate

The title compound is obtained using the procedure described in Example 2 replacing propionic anhydride with chloroacetic anhydride.

15 Melting point : 223-225°C.

## PHARMACOLOGICAL STUDY OF COMPOUNDS OF THE INVENTION

### **EXAMPLE A : Acute toxicity study**

Acute toxicity was evaluated after oral administration to groups each comprising 8 mice (26 ± 2 grams). The animals were observed at regular intervals during the course of the first day, and daily for the two weeks following treatment. The LD<sub>50</sub> (dose that causes the death of 50 % of the animals) was evaluated and demonstrated the low toxicity of the compounds of the invention.

### **EXAMPLE B : Morris water maze test in the mouse**

The anti-amnesic effects of the compounds of the present invention have been evaluated using the Morris water maze test (Morris *et al.*, Nature, 1986, 319, 774-776) in the mouse and scopolamine as amnesic agent. Kunming strain mice (18-24g, Shanghai Experimental Animal Centre) of either sex were used. Mice were placed on the water maze (80x50x20 cm) and trained to find the platform. Following the period of one day's habituation, each mouse received 3 daily training sessions for seven days. Mice were trained to a criterion of finding the platform within 20 seconds and with < 2 errors of entering a dead-end. Once a mouse met the criterion, training was reduced to one daily session until all mice met the criterion. Trained mice were randomly assigned to sub-groups. Compounds under study were dissolved in distilled water and administered by the oral route 40 minutes before behavioural testing. Scopolamine (5 mg/kg, i.p.) was injected 30 minutes before the test. The number of errors and the time for reaching the platform were recorded. Data were expressed as means ± s.e.m. Statistical analysis was performed using ANOVA followed by Duncan's multiple-range test.

Results demonstrate that compounds of the present invention were capable of counteracting in a dose-dependent manner (from 20 to 100 mg/kg) scopolamine-induced memory impairments in the Morris water maze test in the mouse, indicating that such compounds possess anti-amnesic properties. As example, compound of Example 4 gave the following results :

- {

Scopolamine : 0

⇒ Latency to find the platform = 15 s
- {

Example 4 : 0
- {

Scopolamine : 3 mg/kg

⇒ Latency to find the platform = 55 s
- {

Example 4 : 0
- 5
 

{

Scopolamine : 3 mg/kg

⇒ Latency to find the platform = 35 s
- {

Example 4 : 20 mg/kg
- {

Scopolamine : 3 mg/kg

⇒ Latency to find the platform = 25 s
- {

Example 4 : 30 mg/kg

### **EXAMPLE C : Social recognition in the Wistar rat**

- 10 Initially described in 1982 by THOR and HOLLOWAY (J. Comp. Physiol., 1982, 96, 1000-1006), the social recognition test has subsequently been proposed by various authors (DANTZER *et al.*, Psychopharmacology, 1987, 91, 363-368 ; PERIO *et al.*, Psychopharmacology, 1989, 97, 262-268) for studying the mnemocognitive effects of new compounds. The test is based on the natural expression of the olfactory memory of the rat
- 15 and its natural tendency to forget, and allows evaluation of memorisation, by recognition of a young congeneric animal, by an adult rat. A young rat (21 days), taken at random, is placed for 5 minutes in the cage housing an adult rat. With the aid of a video device, the experimenter observes the social recognition behaviour of the adult rat and measures its overall duration. The young rat is then removed from the adult rat's cage and is placed in
- 20 its own cage until the second introduction. The adult rat is given the compound under test and, after 2 hours, is again brought into the presence (5 minutes) of the young rat. The social recognition behaviour is then observed again and its duration measured. The assessment criterion is the difference ( $T_2-T_1$ ), expressed in seconds, between the "recognition" times of the 2 encounters.
- 25 The results obtained show a difference ( $T_2-T_1$ ) ranging from (–10) s to (–36) s for doses ranging from 3 to 30 mg/kg, which shows that the compounds of the invention very greatly enhance memorisation. As example, compound of Example 13 at a dose of 20 mg/kg showed a difference ( $T_2-T_1$ ) of – 36 s, and compound of Example 5 a ( $T_2-T_1$ ) of – 31 s.

**EXAMPLE D : Object recognition in the Wistar rat**

The object recognition test in the Wistar rat was initially developed by ENNACEUR and DELACOUR (Behav. Brain Res., 1988, 31, 47-59). The test is based on the spontaneous exploratory activity of the animal and has the characteristics of episodic memory in humans. This memory test is sensitive to ageing (SCALI *et al.*, Eur. J. Pharmacol., 1997, 325, 173-180) and to cholinergic dysfunctions (BARTOLINI *et al.*, Pharm. Biochem. Behav. 1996, 53(2), 277-283) and is based on the differences in the exploration of 2 objects of fairly similar shape – one familiar, the other new. Prior to the test, the animals are habituated to the environment (an enclosure without an object). In the course of a first session, the rats are placed (3 minutes) in the enclosure, in which there are 2 identical objects. The duration of exploration is measured for each object. In the course of the second session (3 minutes), 24 hours later, 1 of the 2 objects is replaced by a new object. The duration of exploration is measured for each object. The assessment criterion is the difference, Delta, expressed in seconds, between the exploration times for the new object and for the familiar object in the course of the second session. The control animals, previously treated with the carrier by the IP route 30 minutes before each session, explore the familiar object and the new object in an identical manner, which indicates that the object introduced earlier has been forgotten. Animals treated with a compound that facilitates mnemocognition preferentially explore the new object, which indicates that the object introduced earlier has been remembered.

The results obtained show a difference, Delta, ranging from 5 to 11 s, for doses ranging from 0.3 to 10 mg/kg, which shows that the compounds of the invention greatly enhance memorisation. As example, compound of Example 4 showed a Delta of 10 s at a dose of 3 mg/kg.

**EXAMPLE E : NANO<sub>2</sub> induced anoxia in mice**

The neuroprotective effects of the compounds of the present invention have been evaluated in mice. Kunming strain mice of either sex were supplied by Shanghai Experimental Animal Center, Chinese Academy of Sciences (Grade clear, Certificate N°005). Mice

weighing 22-28 g were kept in a 12 hours-light-dark cycle and given food and water *ad libitum*. Compounds under study were dissolved in a 5 % polysorbate-80 solution and orally administered (50 mg/kg) 60 minutes prior to the administration of NaNO<sub>2</sub> at a dose of 225 mg/kg *ip*. Lethality was observed and the prolongation of survival was recorded.

- 5 The results obtained indicate that the compounds of the present invention were able to increase the survival of mice after an *ip* administration of NaNO<sub>2</sub>. These results demonstrate that the compounds of the present invention possess patent anti-anoxic and neuroprotective effects in the mouse. As example, compound of Example 5 shows a prolongation of survival of 31 % at 70 mg/kg *p.o.*.

#### 10 **EXAMPLE F :    Pharmaceutical composition**

Formula for the preparation of 1000 tablets each containing 10 mg of active ingredient:

	(9 $\alpha$ ,13 $\alpha$ )-1-Bromo-3,7-dimethoxy-17-methyl-7,8-didehydromorphinan-4,6-diol .....	10 g
	hydroxypropylcellulose .....	2 g
15	wheat starch .....	10 g
	lactose .....	100 g
	magnesium stearate.....	3 g
	talc .....	3 g